



Iranian Journal of Physics Research, Vol. 20, No. 2, 2020

Geometric phase of a two-level atom near a dissipative and dispersive dielectric slab

S Mohamadi Abdhvand¹, E Amooghorban^{1,2} and AMahdifar^{3,4}

1. Department of Physics, Faculty of Basic Sciences, Shahrekord University, Shahrekord, Iran
2. Nanotechnology Research Center, Shahrekord University, Shahrekord, Iran
3. Department of Physics, Faculty of Physics, University of Isfahan, Isfahan, Iran
4. Quantum Optics Group, Faculty of Physics, University of Isfahan, Isfahan, Iran

E-mail: ehsan.amooghorban@sku.ac.ir

(Received 30 September 2019 ; in final form 14 February 2020)

Abstract

In this paper, we study the geometric phase of a two-level atom near a dielectric slab. For this purpose, by applying the Von-Neumann equation, we obtain the eigenvalues and eigenvectors of the reduced density operator of the atomic system. Then, we obtain the Lamb shift and transition rates of the atomic system in terms of the electromagnetic Green tensor. Finally, by calculating the electromagnetic Green tensor of the system and making use of the kinematic approach, we study the geometric phase of the atomic system near the dielectric slab. We show that the geometric phase can be used as a sensitive probe to the surface-phonon polariton waves.

Keywords: geometric phase, surface polariton-phonon, green tensor, lamb shift and transition rates

For full article, refer to the Persian section