



Iranian Journal of Physics Research, Vol. 21, No. 1, 2021  
DOI:

## Dissipative diffraction in time and early-arrival in continuous quantum-classical transition

S V Mousavi

Department of Physics, Faculty of Sciences, University of Qom, Qom, Iran

E-mail: vmousavi@qom.ac.ir

(Received 15 December 2019 ; in final form 1 December 2020)

### Abstract

Diffraction in time and early-arrival phenomena will be studied in the context of dissipative systems. This theoretical consideration is done by using the linear scaled wave equation proposed in the framework of Caldirola-Kanai theory for the continuous quantum-classical transition. Two simple but physically important and interesting examples are studied, sudden release from a shutter where transient behaviors (diffraction in time) are seen; and also transmission through a time-dependent parabolic barrier where early-arrival is seen. Calculations show that diffraction in time is a non-classical effect and temporary behaviors are gradually suppressed as the friction increases. Furthermore, early-arrival is seen even in the classical regime. This means that early-arrival is not a non-classical phenomenon. This behavior will be explained by means of scaled trajectories which are just Bohmian trajectories in the quantum regime.

**Keywords:** dissipation, Caldirola-Kanai equation, quantum-classical transition, scaled wave equation, diffraction in time, early-arrival, scaled trajectories

For full article, refer to the Persian section