



Iranian Journal of Physics Research, Vol. 23, No. 1, 2022
DOI: 10.47176/ijpr.22.1.61265

Criticality of thermodynamic curvatures for $D \rightarrow 4$ consistent Einstein-Gauss-Bonnet black holes

S A Hosseini Mansoori* and M Rafiee

Faculty of Physics, Shahrood University of Technology, Shahrood, Iran

E-mail: shosseini@shahroodut.ac.ir

(Received 6 June 2021 ; in final form 29 July 2021)

Abstract

In this paper, using Arnowitt-Deser-Misner (ADM) decomposition formalism, we first obtain a neutral black hole solution for consistent $D \rightarrow 4$ Gauss-Bonnet gravity. Then, we study its thermodynamic properties near the critical point by employing new formalism of thermodynamic geometry (NTG). More precisely, critical exponents of the number density difference Δn , the isothermal compressibility $\kappa_T P_c$, the normalized intrinsic R_N , and the normalized extrinsic K_N curvatures are calculated near the critical point for the small/large black holes phase. Our findings show that the critical amplitude of the normalized thermodynamic curvatures are independent of the Gauss-Bonnet coupling values that indicates their universal feature.

Keywords: Gauss-Bonnet black hole, ADM decomposition, intrinsic curvatures and extrinsic curvature

For full article, refer to the Persian section