



Iranian Journal of Physics Research, Vol. 22, No. 4, 2023
DOI: 10.47176/ijpr.22.4.51450

Theoretical investigation of using armchair and zigzag carbon nano rings for DNA sequencing based on density functional theory

E Khodaparast Siyahmazgi, M Qasemnazhand, and F Marsusi*

Department of Physics, Amirkabir university of technology, Tehran, Iran

E-mail: marsusi@aut.ac.ir

(Received 4 May 2022 ; in final form 5 November 2022)

Abstract

Based on density functional theory (DFT) at the B3LYP level, we investigated the interaction of DNA nucleobases with carbon nano-rings in armchair and zigzag shapes. Van der Waals correction was applied to describe the long range term of bipolar interaction. Results indicate that a net electric charge was not transferred between the DNA bases and the carbon nano-rings. This indicates the interaction is of physical type. Outcomes show the following order for the strength of the interaction between the carbon nano-ring (9,9) and the four DNA nucleobases: guanine > adenine > cytosine > thymine. The corresponding order for the zigzag carbon nano-ring (15.0) is adenine \approx guanine > cytosine > thymine, suggesting carbon nano-ring (9,9) may have a potential to specify the sequencing of DNA.

Keywords: DNA sequencing, carbon nano ring, density functional theory, binding energy, Van-der-Waals correction

For full article, refer to the Persian section