



Iranian Journal of Physics Research, Vol. 24, No. 4, 2025
DOI: 10.47176/ijpr.24.4.01995

Investigation of inverse bremsstrahlung absorption in unmagnetized and inhomogeneous plasmas

N Firouzi Farrashbandi^{1*} and A Sid²

1. Department of Physics Education, Farhangian University, P.O. Box 14665-889, Tehran, Iran
2. Laboratory of Physics of Radiations and their Interaction with Matter (PRIMALAB), Department of Physics, Faculty of Matter Sciences, University of Batna 1 - Batna, Algeria

E-mail: na.firouzi@cfu.ac.ir

(Received 26 October 2024 ; in final form 1 December 2024)

Abstract

One of the ways to absorb laser energy by plasma in thermonuclear fusion is called inverse bremsstrahlung absorption. In this process, energy is deposited through inverse bremsstrahlung radiation by plasma electrons. In the current work, inverse bremsstrahlung absorption in unmagnetized and inhomogeneous plasma is calculated. For this purpose, kinetic theory and Fokker-Planck equations are used and the Maxwellian distribution function is considered as the first isotropic distribution function. Then a numerical solution for solving the laser electric field and the plasma dispersion function is proposed. After determining the inverse bremsstrahlung absorption, the effect of laser wavelength and plasma electron temperature on the absorption rate was studied. The results show that increasing the electron temperature and decreasing the laser wavelength increases the amount of absorption in unmagnetized and inhomogeneous plasma. Finally, The difference in inverse bremsstrahlung absorption in homogeneous and inhomogeneous plasmas was investigated.

Keywords: thermonuclear fusion, inverse bremsstrahlung absorption, plasma dispersion function, Fokker-Planck

For full article, refer to the Persian section